

GRADED 3-CALABI-YAU ALGEBRAS AS ORE EXTENSIONS OF 2-CALABI-YAU ALGEBRAS

JI-WEI HE, FRED VAN OYSTAEYEN AND YINHUO ZHANG

ABSTRACT. We study a class of graded algebras obtained from Ore extensions of graded Calabi-Yau algebras of dimension 2. It is proved that these algebras are graded Calabi-Yau and graded coherent. The superpotentials associated to these graded Calabi-Yau algebras are also constructed.

INTRODUCTION

Recently, Smith studied in [Sm2] a remarkable graded Calabi-Yau algebra B of dimension 3 constructed from the octonions. Amongst other things, Smith proved that B is a graded Ore extension of an Artin-Schelter regular algebra of global dimension 2 and uses that fact to show that B is graded 3-Calabi-Yau and graded coherent.

In this note, we show that the Calabi-Yau property and the coherence of B do not occur incidentally. A large class of graded algebras that are Ore extensions of graded Calabi-Yau algebras are themselves graded Calabi-Yau. The main result of this note is the following.

Theorem 0.1. *Let V be a finite dimension vector space with basis $\{x_1, \dots, x_n\}$, let M be an invertible $n \times n$ anti-symmetric matrix, and define*

$$r = (x_1, \dots, x_n)M \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in T(V).$$

Let $A = T(V)/\langle r \rangle$, where $\langle r \rangle$ is the ideal of $T(V)$ generated by r . Let δ be a degree-one graded derivation of $T(V)$ such that $\delta(r) = 0$. Then δ induces a graded derivation $\bar{\delta}$ on A . Let $B = A[z; \bar{\delta}]$ be the Ore extension of A defined by $\bar{\delta}$. Then the following hold:

(i) B is a graded 3-Calabi-Yau algebra.

(ii) Let $\hat{V} = V \oplus \mathbb{k}z$, and $Q = \begin{pmatrix} -1 & 0 \\ 0 & M \end{pmatrix}$. Let $w = (z, x_1, \dots, x_n)Q \begin{pmatrix} r \\ r_1 \\ \vdots \\ r_n \end{pmatrix}$, where

$r_i = z \otimes x_i - x_i \otimes z - \delta(x_i) \in \hat{V} \otimes \hat{V}$ for all $i = 1, \dots, n$. Then $(\alpha \otimes 1 \otimes 1)(w) = (1 \otimes 1 \otimes \alpha)(w)$ for all $\alpha \in (\hat{V})^*$, and $A[z; \bar{\delta}] \cong T(\hat{V})/\langle \partial_{x_i}(w) : i = 0, \dots, n \rangle$, where we set $x_0 = z$ and $\partial_{x_i}(w)$ is the cyclic partial derivative of w with respect to x_i .

2000 *Mathematics Subject Classification.* 16S37, 16S38, 16E65.

Key words and phrases. Calabi-Yau algebra, Ore extension, superpotential.

- (iii) Write $\delta(x_i) = \sum_{s,t=1}^n k_{st}^i x_i \otimes x_j$ for all $i = 1, \dots, n$. Assume there is an integer j such that $k_{jj}^i = 0$ for all $i = 1, \dots, n$, and M is a standard anti-symmetric matrix. Then B is graded coherent.

Most of this note is devoted to the proof of Theorem 0.1. However, we will go a bit further to discuss the properties of the algebra B . Smith's algebra in [Sm2] is an example satisfying the conditions in the theorem. We will provide a few more examples. We remark that any quadratic algebra A defined by an invertible anti-symmetric matrix as in the above theorem is isomorphic to a quadratic algebra defined by a standard anti-symmetric matrix (see Convention 2.3 for the definition), because, for every invertible anti-symmetric matrix M , there is an invertible matrix P such that $P^t M P$ is a standard anti-symmetric matrix.

Remark 0.2. Let V be a finite dimensional vector space with basis $\{x_1, \dots, x_n\}$. Take an element $r \in V \otimes V$. Since $V \otimes V \cong \text{Hom}_{\mathbb{k}}(V^*, V)$, the element r corresponds to a linear map $f_r : V^* \rightarrow V$. The *rank* of r , denoted by $\text{rank}(r)$, is defined to be the rank of f_r (cf. [Z2, Introduction]). One sees

$$\text{rank}(r) = \min\{m \mid r = u_1 \otimes v_1 + \dots + u_m \otimes v_m, \text{ for some } u_i, v_i \in V\}.$$

It has been shown that certain features of the algebra $T(V)/\langle r \rangle$ entirely depend on $\text{rank}(r)$ (cf.

[Z2, Theorem 0.1]). If M is an $n \times n$ matrix and $r = (x_1, \dots, x_n)M \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \in V \otimes V$, then

$\text{rank}(r) = \text{rank}(M)$. Therefore, the condition that M is invertible in Theorem 0.1 is equivalent to the condition that $\text{rank}(r) = n$.

Throughout \mathbb{k} is a fixed field. The unadorned \otimes means $\otimes_{\mathbb{k}}$. Let $U = \bigoplus_{n \in \mathbb{Z}} U_n$ be a graded vector space, and l an integer. We write $U(l)$ for the graded vector space with degree k component $U(l)_k = U_{k+l}$.

A connected graded algebra A is called a *graded Calabi-Yau algebra* of dimension d , or simply *graded d -CY algebra* (cf. [Gin]), if

- (i) A is homologically smooth; that is, A has a finite resolution by finitely generated graded projective left A^e -modules, where $A^e = A \otimes A^{op}$ is the enveloping algebra of A ;
- (ii) the projective dimension of A as a left A^e -module is d , and $\text{Ext}_{A^e}^i(A, A \otimes A) = 0$ if $i \neq d$ and $\text{Ext}_{A^e}^d(A, A \otimes A) \cong A(l)$ for some integer l as a right A^e -module.

We refer to [Z1] (also, cf. [Be] and [DV]) for the basic properties of a graded 2-CY algebra.

1. ORE EXTENSIONS OF GRADED CALABI-YAU ALGEBRAS OF DIMENSION 2

Let V be a vector space with basis $\{x_1, \dots, x_n\}$. Let A be a graded quotient algebra of $T(V)$. If A is a graded 2-CY algebra, then it is defined by an $n \times n$ invertible anti-symmetric matrix M [Z1] (also, cf. [Be, Proposition 3.4]); that is, $A \cong T(V)/\langle r \rangle$ with $r = (x_1, \dots, x_n)M(x_1, \dots, x_n)^t$. Henceforth, we assume $A = T(V)/\langle r \rangle$ with $r = (x_1, \dots, x_n)M(x_1, \dots, x_n)^t$ for some fixed anti-symmetric matrix M . Let $\pi : T(V) \rightarrow A$ be the natural projection map. Since $\text{degree}(r) = 2$, we can, and we will, identify V with A_1 through the projection π .

Let $\delta : V \rightarrow V \otimes V$ be a linear map. Then δ extends in a unique way to a degree-one derivation (also denoted by δ) of $T(V)$. If $\delta(r) \in \langle r \rangle$, then δ induces a derivation $\bar{\delta}$ on A .

From now on, we assume that $\delta(r) \in \langle r \rangle$. Let $B = A[z; \bar{\delta}]$ be the graded Ore extension of A by the derivation $\bar{\delta}$; that is, we view z as an element of degree 1, and $za = az + \bar{\delta}(a)$ for all $a \in A$.

Zhang proved in [Z1] that A is a Koszul algebra of global dimension 2, and the minimal projective resolution of ${}_A \mathbb{k}$ can be written as follows:

$$(1) \quad 0 \longrightarrow A \otimes \mathbb{k}r \xrightarrow{\bar{d}^{-2}} A \otimes V \xrightarrow{\bar{d}^{-1}} A \xrightarrow{\varepsilon} {}_A \mathbb{k} \longrightarrow 0,$$

where ε is the augmentation map, $\bar{d}^{-1}(1 \otimes x) = \pi(x)$ for all $x \in V$, and $\bar{d}^{-2}(1 \otimes r) = r \in A_1 \otimes V$. Since B is an Ore extension of A , B is a Koszul algebra of global dimension 3. Note that B_A is a free A -module. Applying $B \otimes_A -$ to the sequence (1), we obtain the exact sequence

$$(2) \quad 0 \longrightarrow B \otimes \mathbb{k}r \xrightarrow{d^{-2}} B \otimes V \xrightarrow{d^{-1}} B \longrightarrow B/BA_{\geq 1} \longrightarrow 0,$$

where the unlabeled map is the natural projection map, $d^{-1}(1 \otimes x) = \pi(x) \in B_1$ for all $x \in V$, and $d^{-2}(1 \otimes r) = r \in B_1 \otimes V$.

Lemma 1.1. *Suppose that $\delta(r) = 0$ and let $B = A[z; \bar{\delta}]$ be as above. We have the following morphism of cochain complexes:*

$$\begin{array}{ccccc} B \otimes \mathbb{k}r & \xrightarrow{d^{-2}} & B \otimes V & \xrightarrow{d^{-1}} & B \\ f^{-2} \downarrow & & f^{-1} \downarrow & & \downarrow f^0 \\ B \otimes \mathbb{k}r & \xrightarrow{d^{-2}} & B \otimes V & \xrightarrow{d^{-1}} & B, \end{array}$$

where the vertical arrows are left B -module morphisms $f^{-2}(1 \otimes r) = z \otimes r$, $f^{-1}(1 \otimes x) = z \otimes x - \delta(x)$ for all $x \in V$, and $f^0(1) = z$.

Proof. We write $r = \sum_{i=1}^n u_i \otimes x_i$ with all $u_i \in V$, and assume $\delta(x_i) = \sum_{j=1}^n y_{ij} \otimes x_j$ for all $i = 1, \dots, n$ with all $y_{ij} \in V$. We prove the commutativity of the left square. The commutativity of the right one is easy. The identity $\delta(r) = 0$ is equivalent to $\sum_{i=1}^n \delta(u_i) \otimes x_i + \sum_{i=1}^n u_i \otimes \delta(x_i) = 0$, which is in turn equivalent to $\sum_{i=1}^n \delta(u_i) \otimes x_i + \sum_{i=1}^n \sum_{j=1}^n u_i y_{ij} \otimes x_j = 0$. Applying the map $\pi \otimes 1 : T(V) \otimes V \rightarrow A \otimes V$ to the last identity, we obtain $\sum_{i=1}^n \bar{\delta}(u_i) \otimes x_i + \sum_{i=1}^n \sum_{j=1}^n u_i y_{ij} \otimes x_j = 0$. Hence

$$(3) \quad \bar{\delta}(u_i) = - \sum_{j=1}^n u_j y_{ji}$$

for all $i = 1, \dots, n$. The following equations hold:

$$\begin{aligned} f^{-1} \circ d^{-2}(1 \otimes r) &= f^{-1}\left(\sum_{i=1}^n u_i \otimes x_i\right) \\ &= \sum_{i=1}^n u_i z \otimes x_i - \sum_{i=1}^n \sum_{j=1}^n u_i y_{ij} \otimes x_j \\ &= \sum_{i=1}^n \left(u_i z - \sum_{j=1}^n u_j y_{ji}\right) \otimes x_i, \end{aligned}$$

and

$$\begin{aligned} d^{-2} \circ f^{-2}(1 \otimes r) &= d^{-2}(z \otimes r) = \sum_{i=1}^n z u_i \otimes x_i \\ &= \sum_{i=1}^n (u_i z + \bar{\delta}(u_i)) \otimes x_i. \end{aligned}$$

By Equation (3), $f^{-1} \circ d^{-2}(1 \otimes r) = d^{-2} \circ f^{-2}(1 \otimes r)$. Hence the left square of the diagram commutes. \square

The mapping cone of the morphism in Lemma 1.1 provides a graded projective resolution of the trivial module ${}_B \mathbb{k}$ (see also, [GS, Ph]).

Lemma 1.2. *Let r and B be the same as in Lemma 1.1. The minimal projective resolution of ${}_B \mathbb{k}$ is as follows:*

$$0 \longrightarrow B \otimes \mathbb{k}r \xrightarrow{\partial^{-3}} B \otimes \mathbb{k}r \oplus B \otimes V \xrightarrow{\partial^{-2}} B \otimes V \oplus B \xrightarrow{\partial^{-1}} B \longrightarrow \mathbb{k} \longrightarrow 0,$$

$$\text{where } \partial^{-3} = \begin{pmatrix} f^{-2} \\ -d^{-2} \end{pmatrix}, \partial^{-2} = \begin{pmatrix} d^{-2} & f^{-1} \\ 0 & -d^{-1} \end{pmatrix}, \text{ and } \partial^{-1} = (d^{-1}, f^0).$$

Let $A^!$ be the quadratic dual of A . As graded vector spaces $A_0^! \cong \mathbb{k}$, $A_1^! \cong V^*$ and $A_2^! \cong \mathbb{k}r^*$, where $r^* \in (\mathbb{k}r)^*$ defined by $r^*(r) = 1$. The multiplication on $A^!$ is given by: $\alpha\beta = (a_1, \dots, b_n)M(b_1, \dots, b_n)^t r^*$, for $\alpha = a_1 x_1^* + \dots + a_n x_n^*$ and $\beta = b_1 x_1^* + \dots + b_n x_n^*$ in V^* (cf. [HVZ2, Section 3]), where $\{x_1^*, \dots, x_n^*\}$ is the basis of V^* dual to the basis $\{x_1, \dots, x_n\}$. Write $E^i(B) := \text{Ext}_B^i({}_B \mathbb{k}, {}_B \mathbb{k})$ and $E(B) := \bigoplus_{i \geq 0} E^i(B)$. Then $E(B)$ is a graded algebra with the degree i component $E^i(B)$. The minimal projective resolution of ${}_B \mathbb{k}$ above implies that, as graded vector spaces,

$$(4) \quad E(B) \cong A^! \oplus A^!(-1).$$

We write an element in $E(B)$ as (α, β) for some $\alpha, \beta \in A^!$, and demote the Yoneda product on $E(B)$ by $(\alpha, \beta) * (\alpha', \beta')$.

Proposition 1.3. *Assume $\delta(r) = 0$. Then $A[z; \bar{\delta}]$ is a 3-CY algebra.*

Proof. By [HVZ1, Proposition 3.3]) in the Koszul case, $B = A[z; \bar{\delta}]$ is Calabi-Yau if and only if $E(B)$ is a graded symmetric algebra. Recall that a finite dimensional graded algebra $E = \bigoplus_{i \geq 0} E^i$ is graded symmetric if there is an integer d and a homogeneous nondegenerate bilinear form $\langle -, - \rangle : E \times E \rightarrow \mathbb{k}(d)$ such that $\langle \alpha\beta, \gamma \rangle = \langle \alpha, \beta\gamma \rangle$ and $\langle \alpha, \beta \rangle = (-1)^{ij} \langle \beta, \alpha \rangle$ for all homogeneous elements $\alpha \in E^i, \beta \in E^j$ and $\gamma \in E^k$. Since the global dimension of B is 3 and $\dim E^3(B) = 1$, $E(B)$ is graded symmetric if and only if, for all elements $\Phi \in E^1(B), \Theta \in E^2(B)$, $\Phi * \Theta = \Theta * \Phi$. Let $\Phi = (\alpha, k)$ with $\alpha \in A_1^! = V^*$ and $k \in \mathbb{k}$, and $\Theta = (r^*, \beta)$ with $\beta \in V^*$. The element Φ induces a B -module morphism $g : B \otimes V \oplus B \rightarrow {}_B \mathbb{k}$ by $g(1 \otimes x, 1) = \alpha(x) + k$ for all $x \in V$, and the element Θ induces a B -module morphism $h : B \otimes \mathbb{k}r \oplus B \otimes V \rightarrow {}_B \mathbb{k}$ by $h(1 \otimes r, 1 \otimes x) = 1 + \beta(x)$ for all $x \in V$. Consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & B \otimes \mathbb{k}r & \xrightarrow{\partial^{-3}} & B \otimes \mathbb{k}r \oplus B \otimes V & \xrightarrow{\partial^{-2}} & B \otimes V \oplus B \xrightarrow{\partial^{-1}} B \longrightarrow \dots \\ & & \downarrow g_2 & & \downarrow g_1 & & \downarrow g_0 \searrow g \\ \dots & \longrightarrow & B \otimes \mathbb{k}r \oplus B \otimes V & \xrightarrow{\partial^{-2}} & B \otimes V \oplus B & \xrightarrow{\partial^{-1}} & B \longrightarrow {}_B \mathbb{k}, \end{array}$$

where the vertical arrows are B -module morphisms defined as follows. As before, we write $r = \sum_{i=1}^n u_i \otimes x_i$ with all $u_i \in V$, and assume $\delta(x_i) = \sum_{j=1}^n y_{ij} \otimes x_j$ for all $i = 1, \dots, n$ with all $y_{ij} \in V$. Then

$$\begin{aligned} g_0(1 \otimes x_j, 1) &= \alpha(x_j)1 + k1; \\ g_1(1 \otimes r, 1 \otimes x_j) &= \left(\sum_{i=1}^n 1 \otimes u_i \alpha(x_i) - 1 \otimes kx_j - \sum_{i=1}^n 1 \otimes y_{ji} \alpha(x_j), \alpha(x_j)1 \right); \\ g_2(1 \otimes r) &= (1 \otimes kr, \sum_{i=1}^n 1 \otimes u_i \alpha(x_i)), \end{aligned}$$

for all $j = 1, \dots, n$. Since $\delta(r) = 0$, it follows that $\sum_{i=1}^n \delta(u_i) \otimes x_i + \sum_{i=1}^n \sum_{j=1}^n u_i \otimes y_{ij} \otimes x_j = 0$.

Applying the linear map $1 \otimes 1 \otimes \alpha$ to this identity, one obtains:

$$(5) \quad \sum_{i=1}^n \delta(u_i) \alpha(x_i) + \sum_{i=1}^n \sum_{j=1}^n u_i \otimes y_{ij} \alpha(x_j) = 0.$$

Using Equation (5) and the following computations:

$$\begin{aligned} g_1 \circ \partial^{-3}(1 \otimes r) &= g_1(z \otimes r, -r) \\ &= \left(\sum_{i=1}^n z \otimes u_i \alpha(x_i) + \sum_{i=1}^n u_i \otimes kx_i + \sum_{i=1}^n \sum_{j=1}^n u_i \otimes y_{ij} \alpha(x_j), - \sum_{i=1}^n u_i \alpha(x_i) \right), \\ \partial^{-2} \circ g_2(1 \otimes r) &= \partial^{-2} \left(1 \otimes kr, 1 \otimes \sum_{i=1}^n u_i \alpha(x_i) \right) \\ &= \left(kr + \sum_{i=1}^n z \otimes u_i \alpha(x_i) - \sum_{i=1}^n \delta(u_i) \alpha(x_i), - \sum_{i=1}^n u_i \alpha(x_i) \right). \end{aligned}$$

we obtain the identity: $g_1 \circ \partial^{-3}(1 \otimes r) = \partial^{-2} \circ g_2(1 \otimes r)$. Hence $g_1 \circ \partial^{-3} = \partial^{-2} \circ g_2$. Similar computations show that the second square in the diagram commutes. The commutativity of the triangle in the diagram is obvious. Thus, we have $h \circ g_2(1 \otimes r) = h(1 \otimes kr, \sum_{i=1}^n 1 \otimes u_i \alpha(x_i)) = k + \sum_{i=1}^n \beta(u_i) \alpha(x_i)$. By the definition of the Yoneda product, we have $\Theta * \Phi = (r^*, \beta) * (\alpha, k) = kr^* + \beta\alpha$, where $\beta\alpha$ is the product in $A^!$. Similarly, we can show that $\Phi * \Theta = kr^* - \alpha\beta$. Now A is Calabi-Yau, hence $A^!$ is graded symmetric; that is, $\alpha\beta = -\beta\alpha$ for all $\alpha, \beta \in A_1^!$. It follows that $\Phi * \Theta = \Theta * \Phi$. Therefore, $B = A[z; \bar{\delta}]$ is Calabi-Yau. \square

The computation in the proof of Proposition 1.3 has given us the formulas of the Yoneda product of $E(B)$.

Corollary 1.4. *As vector spaces, $E(B) \cong A^! \oplus A^!(-1)$. The Yoneda product of $E(B)$ is given as follows: for $\alpha, \beta \in A_1^!$ and $k, k' \in \mathbb{k}$,*

$$(r^*, \beta) * (\alpha, k) = (\alpha, k) * (r^*, \beta) = kr^* + \beta\alpha,$$

and

$$(\beta, k') * (\alpha, k) = (\beta\alpha, k'\alpha - k\beta - (\beta \otimes \alpha) \circ \delta),$$

where r^* is the basis of $A_2^!$ such that $r^*(r) = 1$.

Proof. The first identity is proved in the proof of Proposition 1.3. Keep the same notions as in the proof of Proposition 1.3. The element (β, k') induces a B -module morphism $g' : B \otimes V \oplus B \rightarrow B\mathbb{k}$ by $g'(1 \otimes x, 1) = \beta(x) + k'$ for all $x \in V$, and $(\beta\alpha, k'\alpha - k\beta - (\beta \otimes \alpha) \circ \delta)$ induces a B -module morphism $f : B \otimes \mathbb{k}r \oplus B \otimes V \rightarrow B\mathbb{k}$ by $f(1 \otimes r, 1 \otimes x_j) = \sum_{i=1}^n \beta(u_i) \alpha(x_i) +$

$k'\alpha(x_j) - k\beta(x_j) - \sum_{i=1}^n \beta(y_{ji})\alpha(x_i)$ for all $j = 1, \dots, n$. By the definition of Yoneda product, $(\beta, k') * (\alpha, k)$ is represented by $g' \circ g_1$. Now $g' \circ g_1(1 \otimes r, 1 \otimes x_j) = \sum_{i=1}^n \beta(u_i)\alpha(x_i) - k\beta(x_j) - \sum_{i=1}^n \beta(y_{ji})\alpha(x_i) + k'\alpha(x_j) = f(1 \otimes r, 1 \otimes x_j)$ for all $j = 1, \dots, n$. Therefore the second identity follows. \square

Let $\epsilon : A^! \rightarrow A^!$ be the automorphism of $A^!$ defined by $\epsilon(\alpha) = -\alpha$ for $\alpha \in A_1^!$ and $\epsilon(\beta) = \beta$ for all $\beta \in A_2^!$. Let ${}_\epsilon A^!$ be the graded $A^!$ -bimodule whose right $A^!$ -action is the regular action, and whose left $A^!$ -action is twisted by the automorphism ϵ ; that is, for all $\gamma, \theta \in A^!$, the left $A^!$ -action $\gamma \cdot \theta = \epsilon(\gamma)\theta$. Let $I = {}_\epsilon A^!(-1)$, and let $E(A^!; I)$ be the trivial extension of $A^!$ by the $A^!$ -bimodule I . By Corollary 1.4, $E(B)$ is isomorphic to $E(A^!; I)$.

Corollary 1.5. *The Yoneda algebra $E(B)$ is isomorphic to the trivial extension of $A^!$ by the $A^!$ -bimodule I .*

Example 1.6. Consider the Calabi-Yau algebra studied by Smith in [Sm2]. Let $\mathbb{k}\langle x_1, \dots, x_6 \rangle$ be the free algebra generated by six elements. Let $A = \mathbb{k}\langle x_1, \dots, x_6 \rangle / \langle r \rangle$, where

$$r = (x_1, \dots, x_6) \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_6 \end{pmatrix}.$$

Define a derivation $\delta : \mathbb{k}\langle x_1, \dots, x_6 \rangle \rightarrow \mathbb{k}\langle x_1, \dots, x_6 \rangle$ by

$$\begin{aligned} \delta(x_1) &= x_4x_2 - x_2x_4 + x_3x_5 - x_5x_3 & \delta(x_2) &= x_1x_4 - x_4x_1 + x_3x_6 - x_6x_3 \\ \delta(x_3) &= x_5x_1 - x_1x_5 + x_6x_2 - x_2x_6 & \delta(x_4) &= x_2x_1 - x_1x_2 + x_5x_6 - x_6x_5 \\ \delta(x_5) &= x_1x_3 - x_3x_1 + x_6x_4 - x_4x_6 & \delta(x_6) &= x_2x_3 - x_3x_2 + x_4x_5 - x_5x_4. \end{aligned}$$

Then $\delta(r) = 0$, and $B = A[z; \bar{\delta}]$ is 3-CY [Sm2].

Keep the assumption that $\delta(r) = 0$. Let $\widehat{V} = V \oplus \mathbb{k}z$. Then $B = A[z; \bar{\delta}]$ is a quotient algebra of $T(\widehat{V})$. Since B is 3-CY, B is defined by a superpotential [Bo, Theorem 3.1]. Let $\{x_1^*, \dots, x_n^*\}$ be the basis of V^* dual to $\{x_1, \dots, x_n\}$. Recall that a *superpotential* is an element $w \in \widehat{V} \otimes \widehat{V} \otimes \widehat{V}$ such that $[\alpha w] = [w\alpha]$ for all $\alpha \in (\widehat{V})^*$, where $[\alpha w] = (\alpha \otimes 1 \otimes 1)(w)$ and $[w\alpha] = (1 \otimes 1 \otimes \alpha)(w)$. Given a superpotential w , the *partial derivative* of w by x_i is defined by $\partial_{x_i}(w) = [x_i^* w]$ (cf. [BSW]). By [Bo, Theorem 3.1], there is a superpotential $w \in \widehat{V} \otimes \widehat{V} \otimes \widehat{V}$ such that $B \cong T(\widehat{V}) / \langle \partial_{x_i}(w) : i = 0, \dots, n \rangle$ where $x_0 = z$. We next show that the superpotential w may be written out explicitly. For $i = 1, \dots, n$, let $r_i = z \otimes x_i - x_i \otimes z - \delta(x_i) \in \widehat{V} \otimes \widehat{V}$. Clearly r, r_1, \dots, r_n are linearly independent in $\widehat{V} \otimes \widehat{V}$, moreover $B \cong T(\widehat{V}) / \langle r, r_1, \dots, r_n \rangle$. Before we construct the general form of the superpotentials, let us look at the following example.

Example 1.7. Let $\mathbb{k}\langle x, y \rangle$ be the free algebra generated by two elements. Let $\delta : \mathbb{k}\langle x, y \rangle \rightarrow \mathbb{k}\langle x, y \rangle$ be a derivation defined by $\delta(x) = bx^2 + cy^2$ and $\delta(y) = ax^2 - bxy - byx$, where $(a, b, c) \in \mathbb{k}^3$. Let $r = xy - yx$. Then it is easy to see that $\delta(r) = 0$. Therefore, δ induces a derivation $\bar{\delta}$ on $A = \mathbb{k}[x, y]$. Now $B = A[z; \bar{\delta}]$ is 3-CY. A straightforward verification shows that $w = yxz + zyx + xzy - xyz - zxy - yzx - ax^3 + cy^3 + bxyx + bx^2y + byx^2$ is a superpotential, and $B \cong \mathbb{k}\langle x, y, z \rangle / \langle \partial_x(w), \partial_y(w), \partial_z(w) \rangle$. Explicitly, the generating relations are $r_1 = zy - yz - ax^2 + bxy + bxy, r_2 = xx - zx + cy^2 + bx^2$ and $r_3 = yx - xy$.

Proposition 1.8. Assume $\delta(r) = 0$. Let $Q = \begin{pmatrix} -1 & 0 \\ 0 & M \end{pmatrix}$, and let $w = (z, x_1, \dots, x_n)Q \begin{pmatrix} r \\ r_1 \\ \vdots \\ r_n \end{pmatrix}$,

where M is an invertible $n \times n$ anti-symmetric matrix, and $r_i = z \otimes x_i - x_i \otimes z - \delta(x_i) \in \widehat{V} \otimes \widehat{V}$ for all $i = 1, \dots, n$. Then

- (i) w is a superpotential;
- (ii) $A[z; \bar{\delta}] \cong T(\widehat{V}) / \langle \partial_{x_i}(w) : i = 0, \dots, n \rangle$, where we set $x_0 = z$.

Proof. Let $\{m_{ij} | i, j = 1, \dots, n\}$ be the entries of M . Then $r = \sum_{i,j=1}^n m_{ij} x_i \otimes x_j$. Since $\delta(r) = 0$, we have $\sum_{i,j=1}^n m_{ij} \delta(x_i) \otimes x_j = - \sum_{i,j=1}^n m_{ij} x_i \otimes \delta(x_j)$. Let us compute the element w .

$$\begin{aligned}
 w &= -z \otimes r + (x_1, \dots, x_n)M \begin{pmatrix} r_1 \\ \vdots \\ r_n \end{pmatrix} \\
 &= - \sum_{i,j=1}^n m_{ij} z \otimes x_i \otimes x_j + \sum_{i,j=1}^n m_{ij} x_i \otimes r_j \\
 &= - \sum_{i,j=1}^n m_{ij} z \otimes x_i \otimes x_j + \sum_{i,j=1}^n m_{ij} x_i \otimes z \otimes x_j \\
 &\quad - \sum_{i,j=1}^n m_{ij} x_i \otimes x_j \otimes z - \sum_{i,j=1}^n m_{ij} x_i \otimes \delta(x_j) \\
 &= - \sum_{i,j=1}^n m_{ij} (z \otimes x_i - x_i \otimes z) \otimes x_j - \sum_{i,j=1}^n m_{ij} x_i \otimes x_j \otimes z + \sum_{i,j=1}^n m_{ij} \delta(x_i) \otimes x_j \\
 &= - \sum_{i,j=1}^n m_{ij} (z \otimes x_i - x_i \otimes z - \delta(x_i)) \otimes x_j - \sum_{i,j=1}^n m_{ij} x_i \otimes x_j \otimes z \\
 &= -r \otimes z + (r_1, \dots, r_n)M^t \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}.
 \end{aligned}$$

Now it is clear that $[x_i^* w] = [w x_i^*]$, and $\partial_{x_i}(w) = r_i$ for all $i = 0, 1, \dots, n$, where $r_0 = r$. \square

2. COHERENCE OF $A[z; \bar{\delta}]$

Notation and notions are as in the previous section. By [Z1, Theorem 0.2], A is Noetherian if and only if $\dim(V) = 2$. Since B is an Ore extension of A in variable z , B/Bz is isomorphic to A as a graded left B -module. Since A is not left Noetherian when $\dim(V) > 2$, neither is B . Similarly, B is not right Noetherian when $\dim(V) > 2$. Summarizing the foregoing argument, we obtain the following property.

Lemma 2.1. $B = A[z; \bar{\delta}]$ is Noetherian if and only if $\dim(V) = 2$.

Piontkovski showed in [Pi, Theorem 4.1] that any connected graded algebra with a single quadratic relation is graded coherent. Hence A is a graded coherent algebra. So, it is natural

to ask whether B is a graded coherent algebra. The answer is affirmative. However, the proof of this property is not trivial because an Ore extension of a coherent ring needs not be coherent. In fact, there is a commutative coherent ring R such that the polynomial extension $R[z]$ is not coherent [So]. Some other results about the coherence of polynomial rings may be found in [GV]. Let us recall the definition of a graded coherent algebra.

A graded algebra D is called a *graded left coherent algebra* if one of the following equivalent conditions is satisfied:

- (i) every finitely generated graded left ideal of D is finitely presented; that is, if I is a graded left ideal of D then there is a finitely graded free D -module F and a surjective morphism $g : F \rightarrow I$ of graded modules such that $\ker g$ is also a finitely generated D -module;
- (ii) every finitely generated graded submodule of a finitely presented graded module is finitely presented;
- (iii) the category of all finitely presented graded left D -modules is an abelian category.

Similarly we can define a *graded right coherent algebra*. If a graded algebra is both graded left and right coherent, then it is called a *graded coherent algebra*.

Let $W = \oplus_{i \geq 0} W_i$ be a graded vector space with $\dim(W_i) < \infty$ for all i . Recall that the Hilbert series of W is defined to be the power series $H_W(t) = \sum_{i \geq 0} \dim(W_i)t^i$.

Lemma 2.2. *Let V be a vector space of dimension $n \geq 4$ with basis $\{x_1, \dots, x_n\}$, and let*

$$(6) \quad M = \begin{pmatrix} 0 & \cdots & 0 & 0 & \cdots & 1 \\ \vdots & & \vdots & \vdots & & \vdots \\ 0 & \cdots & 0 & 1 & \cdots & 0 \\ 0 & \cdots & -1 & 0 & \cdots & 0 \\ \vdots & & \vdots & \vdots & & \vdots \\ -1 & \cdots & 0 & 0 & \cdots & 0 \end{pmatrix}$$

be the invertible $n \times n$ anti-symmetric matrix with entries in the anti-diagonal line 1 or -1 and others 0. Let $r = (x_1, \dots, x_n)M(x_1, \dots, x_n)^t$, and $A = T(V)/\langle r \rangle$. Let δ be a derivation on $T(V)$ of degree 1. We write $\delta(x_i) = \sum_{s,t=1}^n k_{st}^i x_s \otimes x_t$ for all $i = 1, \dots, n$. Assume that $k_{nn}^i = 0$ for all $i = 1, \dots, n$ and $\delta(r) = 0$. Let $\bar{\delta}$ be the derivation on A induced by δ . Write $B = A[z; \bar{\delta}]$. Then the following hold.

- (i) *Let I be the ideal of B generated by the elements x_1, \dots, x_{n-1} . Then $B/I \cong \mathbb{k}[X, Z]$, where $\mathbb{k}[X, Z]$ is the commutative polynomial algebra in variables X and Z ;*
- (ii) *Let $L = \mathbb{k}x_1 \oplus \cdots \oplus \mathbb{k}x_{n-1}$ and $L' = \mathbb{k}x_2 \oplus \cdots \oplus \mathbb{k}x_{n-1}$. Then, as left B -modules, $I \cong B \otimes (L \oplus L'x_n \oplus L'x_n^2 \oplus \cdots)$, where $L'x_n^k$ ($k \geq 1$) is the vector space spanned by the elements $x_2x_n^k, \dots, x_{n-1}x_n^k$.*

Convention 2.3. We call an $n \times n$ ($n \geq 2$) invertible anti-symmetric matrix of the form (6) a *standard anti-symmetric matrix*. If M is an invertible anti-symmetric matrix, there is an invertible matrix P such that $P^t M P$ is standard.

Proof of Lemma 2.2. (i) By assumption, $\delta(x_n) = \sum_{s,t=1}^n k_{st}^n x_s \otimes x_t$ and $k_{nn}^n = 0$. Therefore $\bar{\delta}(x_n) \in I$ and B/I is a commutative algebra. There is an algebra morphism $g : \mathbb{k}[X, Z] \rightarrow B/I$ defined by $g(X) = x_n$ and $g(Z) = z$. Next, we want to construct an algebra morphism from B/I to $\mathbb{k}[X, Z]$. As before, write $\hat{V} = V \oplus \mathbb{k}z$. Firstly, we define $f : T(\hat{V}) \rightarrow \mathbb{k}[X, Z]$ by

letting $f(x_i) = 0$ for all $i = 1, \dots, n-1$, $f(x_n) = X$ and $f(z) = Z$. Denote by $\langle x_1, \dots, x_{n-1} \rangle$ and by $\langle z \otimes x_n - x_n \otimes z \rangle$ the ideals of $T(\widehat{V})$ respectively generated by x_1, \dots, x_{n-1} and by $z \otimes x_n - x_n \otimes z$. Obviously, $\langle x_1, \dots, x_{n-1} \rangle + \langle z \otimes x_n - x_n \otimes z \rangle \subseteq \ker f$. Recall that B is a Koszul algebra and $B = T(\widehat{V})/J$ where $J = \langle r, z \otimes x_1 - x_1 \otimes z - \delta(x_1), \dots, z \otimes x_n - x_n \otimes z - \delta(x_n) \rangle$. Since $\delta(x_i) = \sum_{s,t=1}^n k_{st}^i x_s \otimes x_t$ such that $k_{nn}^i = 0$ for all $i = 1, \dots, n$, it follows that $\delta(x_i) \in \langle x_1, \dots, x_{n-1} \rangle$ for all $i = 1, \dots, n$. Hence $r, z \otimes x_1 - x_1 \otimes z - \delta(x_1), \dots, z \otimes x_{n-1} - x_{n-1} \otimes z - \delta(x_{n-1}) \in \langle x_1, \dots, x_{n-1} \rangle \subseteq \ker f$. Now $z \otimes x_n - x_n \otimes z - \delta(x_n) \in \langle z \otimes x_n - x_n \otimes z \rangle + \langle x_1, \dots, x_{n-1} \rangle \subseteq \ker f$. Hence $J \subseteq \ker f$. Therefore, f induces an algebra morphism $\bar{f} : B \rightarrow \mathbb{k}[X, Z]$. Obviously, $\ker \bar{f} \supseteq I$. Hence \bar{f} in turn induces an algebra morphism $\hat{f} : B/I \rightarrow \mathbb{k}[X, Z]$. Now it is easy to see that $\hat{f} \circ g = id = g \circ \hat{f}$. The statement (i) follows.

(ii) Here we make use of the technique from [Sm2, Proposition 7.3]. Let $\mu : B \otimes B \rightarrow B$ be the multiplication of B . Then the restriction of μ defines a left B -module morphism (also denoted by μ):

$$\mu : B \otimes (L \oplus L'x_n \oplus L'x_n^2 \oplus \dots) \rightarrow I.$$

We claim that μ is surjective. In fact, if we can show that the image $I' = \text{im}(\mu)$ is also an ideal of B , then $I = I'$. So, it suffices to show that $I'x_n \subseteq I'$ and $I'z \subseteq I'$. Following the generating relation of A , we have $x_1x_n = x_nx_1 + (x_2x_{n-1} - x_{n-1}x_2) + \dots + (x_{\frac{n}{2}}x_{\frac{n}{2}+1} - x_{\frac{n}{2}+1}x_{\frac{n}{2}}) \in BL \subseteq I'$. Therefore $I'x_n \subseteq I'$. In particular, $\bar{\delta}(x_i) \in I'$ for all $i = 1, \dots, n$. On the other hand, since $x_iz = zx_i - \delta(x_i)$, it follows that $x_iz \in I'$ for all $i = 1, \dots, n-1$. For $2 \leq i \leq n-1$, we have $x_ix_nz = x_i(zx_n - \bar{\delta}(x_n)) = x_izx_n - x_i\bar{\delta}(x_n) \in I'x_n + x_iI' \subseteq I'$. Now assume $x_ix_n^jz \in I'$ for all $j < k$ and $2 \leq i \leq n-1$. Then

$$x_ix_n^kz = x_ix_n^{k-1}(zx_n - \bar{\delta}(x_n)) = (x_ix_n^{k-1}z)x_n - x_ix_n^{k-1}\bar{\delta}(x_n) \in I'x_n + x_ix_n^{k-1}I' \subseteq I'.$$

Hence $I'z \subseteq I'$. The claim follows. To show that μ is injective, we only need to compare the Hilbert series of I and that of $F := B \otimes (L \oplus L'x_n \oplus L'x_n^2 \oplus \dots)$. Write $W = L \oplus L'x_n \oplus L'x_n^2 \oplus \dots$. Clearly $H_F(t) = H_B(t) \cdot H_W(t)$. We have

$$H_W(t) = (n-1)t + (n-2)t^2 + (n-2)t^3 + \dots = ((n-1)t - t^2)(1-t)^{-1}.$$

The exact sequence $0 \rightarrow I \rightarrow B \rightarrow B/I \rightarrow 0$ implies $H_I(t) = H_B(t) - H_{B/I}(t)$. Since B is Koszul of global dimension 3, it follows that $H_B(t) = (1 - (n+1)t + (n+1)t^2 - t^3)^{-1}$ by [Sm1, Theorem 5.9] and the isomorphism (4) of the previous section. By (i), $H_{B/I}(t) = (1-t)^{-2}$. Hence

$$\begin{aligned} H_I(t) &= (1 - (n+1)t + (n+1)t^2 - t^3)^{-1} - (1-t)^{-2} \\ &= (1 - (n+1)t + (n+1)t^2 - t^3)^{-1} \cdot ((n-1)t - t^2)(1-t)^{-1} \\ &= H_B(t) \cdot H_W(t) \\ &= H_F(t). \end{aligned}$$

Therefore μ is injective. So, (ii) follows. \square

Proof of the statement (iii) of Theorem 0.1. If $n = 2$, then $A = \mathbb{k}[x_1, x_2]$. We obtain that $B = A[z; \bar{\delta}]$ is Noetherian, and hence coherent. Now assume $n \geq 4$. We only prove the statement when $j = n$ in the assumption, that is, $k_{nn}^i = 0$ for all $i = 1, \dots, n$. When $j \neq n$, the statement can be proved similarly. By Lemma 2.2, there is an exact sequence $0 \rightarrow I \rightarrow B \rightarrow B/I \rightarrow 0$ such that B/I is a polynomial algebra in two variables and I is a free graded left B -module. By [Pi, Proposition 3.2], B is graded right coherent. Note that the left version of Lemma 2.2(ii) holds too. Hence B is also graded left coherent. \square

As a special case of the statement (iii) of Theorem 0.1, we have the following result, which can be viewed as a noncommutative version of [GV, Theorem 4.3].

Proposition 2.4. *Let A be a connected graded 2-CY algebra. Then $A[z]$ is a graded coherent algebra.*

Proof. By [Z1, Theorem 0.1] (also, cf. [Be, Proposition 3.4]), A is defined by an invertible anti-symmetric matrix M , that is, $A = T(V)/\langle r \rangle$ with $r = (x_1, \dots, x_n)M(x_1, \dots, x_n)^t$. For an invertible anti-symmetric matrix M , there is an invertible matrix P such that $P^t M P$ is a standard invertible anti-symmetric matrix. Then the algebras defined by M and $P^t M P$ respectively are isomorphic to each other. Hence we may assume that the anti-symmetric matrix M itself is standard. Now by (iii) of Theorem 0.1, we see that $A[z]$ is graded coherent. \square

Now assume that $B = A[z; \bar{\delta}]$ is graded coherent. We may form a noncommutative projective space from B . Following [Po], we denote by $\text{coh}B$ the category of all finitely presented graded left B -modules, and by $\text{fdim}B$ the category of all finite dimensional graded left B -modules. Since B is graded coherent, $\text{fdim}B$ is a Serre subcategory of $\text{coh}B$. Hence the quotient category

$$\text{cohproj}B := \text{coh}B / \text{fdim}B$$

is also an abelian category. Since B is Koszul and 3-CY, B is Artin-Schelter regular with Gorenstein parameter -3 . Hence the Beilinson algebra of B (for the terminology, see [MM, Definition 4.7]) is

$$\nabla B = \begin{pmatrix} \mathbb{k} & B_1 & B_2 \\ 0 & \mathbb{k} & B_1 \\ 0 & 0 & \mathbb{k} \end{pmatrix}.$$

Let $\text{mod}\nabla B$ be the category of finite dimensional left ∇B -modules. Then by [MM, Theorem 4.14], we have the following corollary.

Corollary 2.5. *If the conditions of Theorem 0.1 are satisfied, then there is an equivalence of triangulated categories:*

$$D^b(\text{cohproj}B) \cong D^b(\text{mod}\nabla B),$$

where $D^b(-)$ is the bounded derived category of the corresponding abelian category.

Acknowledgement. The authors are very grateful to the referee for his/her careful reading of the manuscript, numerous comments and suggestions, which has greatly improved the paper. In particular, Remark 0.2 was suggested to the authors by the referee. The work is supported by an FWO-grant and grants from NSFC (No. 11171067) and NSF of Zhejiang Province (No. LY12A01013). This work is also in part supported by SRF for ROCS, SEM.

REFERENCES

- [Be] R. Berger, Gerasimov's theorem and N-Koszul algebras, J. London Math. Soc. 79 (2009), 631–648.
- [Bo] R. Bocklandt, Graded Calabi-Yau algebras of dimension 3, J. Pure Appl. Algebra 212 (2008), 14–32.
- [BSW] R. Bocklandt, T. Schedler, M. Wemyss, Superpotentials and higher order derivations, J. Pure Appl. Algebra 214 (2010), 1501–1522.
- [CS] T. Cassidy, B. Shelton, Generalizing the notion of Koszul algebra, Math. Z. 260 (2008), 93–114.
- [DV] M. Dubois-Violette, Multilinear forms and graded algebras, J. Algebra 317 (2007), 198–225.
- [Gin] V. Ginzburg, Calabi-Yau algebras, arXiv:math/0612139.
- [GS] N. S. Gopalakrishnan, R. Sridharan, Homological dimension of Ore-extensions, Pacific J. Math. 19 (1966), 67–75.

- [GV] B. V. Greenberg, W. V. Vasconcelos, Coherence of polynomial rings, Proc. Amer. Math. Soc. 54 (1976), 59–64.
- [HVZ1] J.-W. He, F. Van Oystaeyen, Y. Zhang, Hopf algebra actions on differential graded algebras and applications, Bull. Belg. Math. Soc. Simon Stevin 18 (2011), 99–111.
- [HVZ2] J.-W. He, F. Van Oystaeyen, Y. Zhang, Deformations of Koszul Artin-Schelter Gorenstein algebras, Manuscripta Math., DOI: 10.1007/s00229-012-0580-z; arXiv:1302.0390.
- [MM] H. Minamoto, I. Mori, Structure of AS-Gorenstein algebras, Adv. Math. 226 (2011), 4061–4095.
- [Ph] C. Phan, The Yoneda algebra of a graded Ore extension, Comm. Algebra 40 (2012), 834–844.
- [Pi] D. Piontkovski, Coherent algebras and noncommutative projective lines, J. Algebra 319 (2008), 3280–3290.
- [Po] A. Polishchuk, Noncommutative proj and coherent algebras, Math. Res. Let. 12 (2005), 63–72.
- [Sm1] S. P. Smith, Some graded algebras related to elliptic curves, CMS Conf. Proc. 19 (1996), Amer. Math. Soc., Providence, RI, 315–348.
- [Sm2] S. P. Smith, A 3-Calabi-Yau algebra with G_2 symmetry constructed from the Octonions, arXiv:1104.3824v1.
- [So] J. Soublin, Un anneau cohérent dont l’anneau des polynômes n’est pas cohérent, C. R. Acad. Sci. Paris Sér. A-B 267 (1968), A241–A243.
- [Z1] J. J. Zhang, Non-Noetherian regular rings of dimension 2, Proc. Amer. Math. Soc. 126 (1998), 1645–1653.
- [Z2] J. J. Zhang, Quadratic algebras with few relations, Glasgow Math. J. 39 (1997), 323–332.

J.-W. HE

DEPARTMENT OF MATHEMATICS, SHAOXING COLLEGE OF ARTS AND SCIENCES, SHAOXING ZHEJIANG 312000, CHINA

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF ANTWERP, MIDDELHEIMLAAN 1, B-2020 ANTWERP, BELGIUM

E-mail address: `jwhe@usx.edu.cn`

F. VAN OYSTAEYEN

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF ANTWERP, MIDDELHEIMLAAN 1, B-2020 ANTWERP, BELGIUM

E-mail address: `fred.vanoystaeyen@ua.ac.be`

Y. ZHANG

DEPARTMENT WNI, UNIVERSITY OF HASSELT, UNIVERSITAIRE CAMPUS, 3590 DIEPENBEEK, BELGIUM

E-mail address: `yinhuo.zhang@uhasselt.be`